Transcriptional distinctions between normal and glioma-derived A2B5[†] progenitor cells identify genes dysregulated at all stages of gliomagenesis

Romane M. Auvergne^{1,2,*}, Fraser J. Sim^{1,2,5,*}, Su Wang^{1,2}, Devin Chandler-Militello^{1,2}, Jaclyn Burch^{1,2}, Yazan Al Fanek^{1,2}, Danielle Davis^{1,2}, Xiaojie Li^{1,2}, Abdellatif Benraiss^{1,2}, Nimish Mohile^{1,2}, Kevin Walter³, Pragathi Achanta⁶, Mahlon Johnson⁴, Alfredo Quinones-Hinojosa⁶, Heide Ford⁷, Sridaran Natesan⁸, Steven A. Goldman^{1,2}

SUPPLEMENTAL FIGURES AND LEGENDS

Figure S1, related to Figure 1

Figure S2, related to Figure 1

Figure S3, related to Figure 1

Figure S4, related to Figure 1

Figure S5, related to Figure 2

Figure S6, related to Figure 2

Figure S7, related to Figure 5

Figure S8, related to Figure 5

Figure S9, related to Figure 6

Figure S10, related to Figure 7

SUPPLEMENTAL TABLES AND LEGENDS

Table S1, related to Figure 1

Table S2, related to Figure 2

Table S3, related to Figure 2. Provided as an Excel file

Table S4, related to Figure 2. Provided as an Excel file

Table S5, related to Figure 2. Provided as an Excel file

Table S6, related to Figure 2. Provided as an Excel file

Table S7, related to Figure 2. Provided as an Excel file

Table S8, related to Figure 2

Table S9, related to Figures 2-4. Provided as an Excel file

Table S10, related to Figure 7

Table S11, related to Figures 1,7

SUPPLEMENTAL EXPERIMENTAL PROCEDURES

SUPPLEMENTAL REFERENCES